

GROUPOID ALGEBRAS AS CUNTZ-PIMSNER ALGEBRAS

ADAM RENNIE, DAVID ROBERTSON, AND AIDAN SIMS

ABSTRACT. We show that if G is a second countable locally compact Hausdorff étale groupoid carrying a suitable cocycle $c : G \rightarrow \mathbb{Z}$, then the reduced C^* -algebra of G can be realised naturally as the Cuntz-Pimsner algebra of a correspondence over the reduced C^* -algebra of the kernel G_0 of c . If the full and reduced C^* -algebras of G_0 coincide, we deduce that the full and reduced C^* -algebras of G coincide. We obtain a six-term exact sequence describing the K -theory of $C_r^*(G)$ in terms of that of $C_r^*(G_0)$.

In this short note we provide a sufficient condition for a groupoid C^* -algebra to have a natural realisation as a Cuntz-Pimsner algebra. The advantage to knowing this is the complementary knowledge one obtains from the two descriptions.

Our main result starts with a second-countable locally compact Hausdorff étale groupoid G with a continuous cocycle into the integers which is unperforated in the sense that G is generated by $c^{-1}(1)$. We then show that the reduced groupoid C^* -algebra $C_r^*(G)$ is the Cuntz-Pimsner algebra of a natural C^* -correspondence over the reduced C^* -algebra $C_r^*(c^{-1}(0))$ of the kernel of c .

We also show that if $C^*(c^{-1}(0))$ and $C_r^*(c^{-1}(0))$ coincide, then $C^*(G)$ and $C_r^*(G)$ coincide as well. We finish by applying results of Katsura to present a six-term exact sequence relating the K -theory of $C_r^*(G)$ to that of $C_r^*(c^{-1}(0))$.

Notation 1. For the duration of the paper, we fix a second-countable locally compact Hausdorff étale groupoid G with unit space $G^{(0)}$ and a continuous cocycle $c : G \rightarrow \mathbb{Z}$. We suppose that c is *unperforated* in the sense that if $c(\gamma) = n > 0$, then there exist composable $\gamma_1, \dots, \gamma_n$ such that each $c(\gamma_i) = 1$ and $\gamma = \gamma_1 \dots \gamma_n$. For $n \in \mathbb{Z}$ we write $G_n := c^{-1}(n)$. Recall from [6] that for $u \in G^{(0)}$, we write $G_u = s^{-1}(u)$ and $G^u = r^{-1}(u)$. The convolution multiplication of functions is denoted by juxtaposition, as is multiplication in the C^* -completions.

Remarks.

- (i) Observe that $G^{(0)} \subseteq G_0$, and since c is continuous, G_0 is a clopen subgroupoid of G .
- (ii) If c is strongly surjective in the sense that $c(r^{-1}(u)) = \mathbb{Z}$ for all $u \in G^{(0)}$ [1, Definition 5.3.7], and $\gamma \in G_n$, then there exists $\alpha \in r^{-1}(r(\gamma)) \cap G_1$, and then $\gamma = \alpha(\alpha^{-1}\gamma) \in G_1 G_{n-1}$. So an induction shows that if c is strongly surjective then it is unperforated.

Lemma 2. *Let $A_0 \subseteq C_r^*(G)$ be the completion of $\{f \in C_c(G) : \text{supp}(f) \subseteq G_0\}$. Then there is an isomorphism $I_0 : C_r^*(G_0) \rightarrow A_0$ extending the identity map on $C_c(G_0)$.*

Proof. Fix $u \in G^{(0)}$, let π_u^0 be the regular representation of $C^*(G_0)$ on $\mathcal{H}_u^0 := \ell^2((G_0)_u)$ and let π_u be the regular representation of $C^*(G)$ on $\mathcal{H}_u := \ell^2(G_u)$. For $a \in C_c(G_0)$, the subspace $\mathcal{H}_u^0 \subseteq \mathcal{H}_u$ is reducing for the operator $\pi_u(a)$. Let $P_0 : \mathcal{H}_u \rightarrow \mathcal{H}_u^0$ be the orthonormal projection. We have

$$\|\pi_u^0(a)\| = \|P_0 \pi_u(a) P_0\| \leq \|\pi_u(a)\|.$$

Taking the supremum over all u , we deduce that $\|a\|_{C_r^*(G_0)} \leq \|a\|_{C_r^*(G)}$. So there is a C^* -homomorphism $\pi : A_0 \rightarrow C_r^*(G_0)$ extending the identity map on $C_c(G_0)$.

2010 *Mathematics Subject Classification.* 46L05.

Key words and phrases. Groupoid; Cuntz-Pimsner; C^* -correspondence; Hilbert module.

Restriction of functions induces a faithful conditional expectation[†] $\Psi : C_r^*(G) \rightarrow C_0(G^{(0)})$ and a faithful conditional expectation $\Psi_0 : C_r^*(G_0) \rightarrow C_0(G_0^{(0)}) = C_0(G^{(0)})$. We clearly have $\pi \circ \Psi = \Psi_0 \circ \pi$, and so a standard argument (see [7, Lemma 3.13]) shows that π is faithful and hence isometric. It follows immediately that $I_0 := \pi^{-1} : C_r^*(G_0) \rightarrow A_0$ is an isometric embedding of $C_r^*(G_0)$ in $C_r^*(G)$ as claimed. $\square$

Lemma 3. *Let $X(G)$ be the completion in $C_r^*(G)$ of the subspace $\{f \in C_c(G) : \text{supp}(f) \subseteq G_1\}$. Then $X(G)$ is a C^* -correspondence over $C_r^*(G_0)$ with module actions $a \cdot \xi = I_0(a)\xi$ and $\xi \cdot a = \xi I_0(a)$ and inner product $\langle \cdot, \cdot \rangle_{C_r^*(G_0)} : X(G) \times X(G) \rightarrow C_r^*(G)$ given by $\langle \xi, \eta \rangle_{C_r^*(G_0)} = I_0^{-1}(\xi^* \eta)$.*

Proof. Since c is a cocycle, we have $G_1^{-1} = G_{-1}$. That c is a cocycle also implies that if $f \in C_c(G_m)$ and $g \in C_c(G_n)$ then $fg \in C_c(G_{m+n})$ and $f^* \in C_c(G_{-m})$, and it follows that $C_c(G_1)C_c(G_0), C_c(G_0)C_c(G_1) \subseteq C_c(G_1)$ and that $C_c(G_1)^*C_c(G_1) \subseteq C_c(G_0)$. The C^* -identity and Lemma 2 show that $\langle \cdot, \cdot \rangle_{C_r^*(G)}$ is isometric in the sense that $\|\langle \xi, \xi \rangle_{C_r^*(G_0)}\| = \|\xi\|_{C_r^*(G)}$, and in particular is positive definite. Continuity therefore guarantees that the operations described determine a Hilbert-module structure on $X(G)$. The left action of $C_r^*(G_0)$ on $X(G)$ is adjointable since $\langle a \cdot \xi, \eta \rangle_{C_r^*(G)} = (a\xi)^* \eta = \xi^* (a^* \eta) = \langle \xi, a^* \cdot \eta \rangle_{C_r^*(G)}$. $\square$

Notation 4. We write I_1 for the inclusion map $X(G) \hookrightarrow C_r^*(G)$.

Lemma 5. *The embeddings $I_1 : X(G) \hookrightarrow C_r^*(G)$ and $I_0 : C_r^*(G_0) \hookrightarrow C_r^*(G)$ form a Toeplitz representation (I_1, I_0) of $X(G)$ in $C_r^*(G)$.*

Proof. For $\xi \in X(G)$ and $a \in C_r^*(G_0)$, we have $I_1(\xi)I_0(a) = \xi a = I_1(\xi \cdot a)$ by definition of the actions on $X(G)$. For $\xi, \eta \in C_c(G_1)$, we have

$$I_0(\langle \xi, \eta \rangle_{A_0}) = I_0(\xi^* \eta) = \xi^* \eta = I_1(\xi)^* I_1(\eta).$$

So (I_1, I_0) is a Toeplitz representation as claimed. $\square$

We aim to show that (I_1, I_0) is in fact a Cuntz-Pimsner covariant representation, but we have a little work to do first.

Lemma 6. *For each integer $n > 0$, the space*

$$\text{span}\{f_1 f_2 \cdots f_n : f_i \in C_c(G_1) \text{ for all } i\}$$

is dense in $C_c(G_n)$, and we have $C_c(G_n)^* = C_c(G_{-n})$.

Proof. Fix $x \neq y \in G_n$. Since $G_n = c^{-1}(n)$ is locally compact, by the Stone-Weierstrass theorem it suffices to construct functions $f_1, \dots, f_n \in C_c(G_1)$ such that $(f_1 \cdots f_n)(x) = 1$ and $(f_1 \cdots f_n)(y) = 0$.

Since c is unperforated, there exist $x_1, \dots, x_n \in G_1$ such that $x = x_1 \cdots x_n$. Choose a precompact open bisection $U_n \subseteq G_1$ such that $x_n \in U_n$ and such that if $s(x) \neq s(y)$ then $s(y) \notin s(U_n)$; this is possible because s is a local homeomorphism. For each $1 \leq i \leq n-1$ choose a precompact open bisection $U_i \subseteq G_1$ with $x_i \in U_i$. Then $U_1 \cdots U_n$ is a precompact open bisection containing x because multiplication in G is continuous and open. If $s(x) = s(y)$, then $y \notin U_1 \cdots U_n$ because the latter is a bisection; and if $s(x) \neq s(y)$ then $y \notin U_1 \cdots U_n$ by choice of U_n .

By Urysohn's Lemma, for each i there exists $f_i \in C_c(U_i)$ with $f_i(x_i) = 1$. Then $(f_1 \cdots f_n)(x) = 1$ by construction, and the convolution formula says that

$$\text{supp}(f_1 \cdots f_n) = \text{supp}(f_1) \text{supp}(f_2) \cdots \text{supp}(f_n) \subseteq U_1 \cdots U_n,$$

which yields $(f_1 \cdots f_n)(y) = 0$.

[†]To see this observe that for $f \in C_c(G)$, we have $f(u) = (\pi_u(f)\delta_u|\delta_u)$, giving $\|f|_{G^{(0)}}\| = \sup_u (\pi_u(f)\delta_u|\delta_u) \leq \sup_u \|\pi_u(f)\| = \|f\|_{C_r^*(G)}$, so restriction induces an idempotent map $\Psi : C_r^*(G) \rightarrow C_0(G^{(0)})$ of norm one. Hence [2, Theorem II.6.10.2] implies that Ψ is a conditional expectation. Given $a \in C_r^*(G) \setminus \{0\}$, [6, Proposition II.4.2] shows that $a \in C_0(G)$, so $a(\gamma) \neq 0$ for some γ . Then $\Psi(a^* a)(s(\gamma)) = \sum_{\alpha \in G_{s(\gamma)}} \overline{a}(\alpha^{-1}) a(\alpha) \geq |a(\gamma)|^2 > 0$, so Ψ is faithful.

The involution formula $f^*(\gamma) = \overline{f}(\gamma^{-1})$ and the cocycle property $c(\gamma^{-1}) = -c(\gamma)$ show that $f \in C_c(G_n)$ if and only if $f^* \in C_c(G_{-n})$. $\square$

Corollary 7. *There is an injection $\psi : \mathcal{K}(X(G)) \rightarrow C_r^*(G_0)$ such that $\psi(\theta_{\xi,\eta}) = \xi\eta^*$ for all $\xi, \eta \in C_c(G_1)$.*

Proof. The discussion on page 202 of [5] (see also [3, Section 1]) shows that there is a homomorphism $I_1^{(1)} : \mathcal{K}(X(G)) \rightarrow C_r^*(G)$ such that $I_1^{(1)}(\theta_{\xi,\eta}) = I_1(\xi)I_1(\eta)^* = \xi\eta^* = I_0(\xi\eta^*)$. Since I_0 is isometric, the composition $\psi := I_0^{-1} \circ I_1^{(1)}$ is a homomorphism satisfying the desired formula, and we need only show that ψ is injective.

If $T \in \mathcal{K}(X(G))$ and $\psi(T) = 0$, then $I_1^{(1)}(T) = 0$ as well. For $\eta, \xi, \zeta \in X(G)$, we have

$$I_1^{(1)}(\theta_{\eta,\xi})I_1(\zeta) = I_1(\eta)I_1(\xi)^*I_1(\zeta) = I_1(\eta \cdot \langle \xi, \zeta \rangle_{A_0}) = I_1(\theta_{\eta,\xi}(\zeta)),$$

and linearity and continuity show that for all $S \in \mathcal{K}(X(G))$ and $\zeta \in X(G)$, we have the formula $I_1^{(1)}(S)I_1(\zeta) = I_1(S\zeta)$. In particular, $0 = I_1^{(1)}(T)I_1(\zeta) = I_1(T\zeta)$ for all $\zeta \in X(G)$. Since I_1 is isometric, we deduce that $T\zeta = 0$ for all $\zeta \in C_c(G_1)$, and so $T = 0$. So ψ is injective. $\square$

Lemma 8. *Let $\phi = \phi_{X(G)} : C_r^*(G_0) \rightarrow \mathcal{L}(X(G))$ be the homomorphism implementing the left action. Then*

- (1) $\ker(\phi) = C_r^*(G_0) \cap C_0(\{g \in G_0 : s(g) \notin r(G_1)\})$;
- (2) $\ker(\phi)^\perp = C_r^*(G_0) \cap C_0(\{g \in G_0 : s(g) \in r(G_1)\}) = \overline{\text{span}}\{fg^* : f, g \in C_c(X_1)\}$;
- (3) $\phi_{X(G)}(a) \in \mathcal{K}(X(G))$ for all $a \in C_r^*(G)$, and the Katsura ideal $J_{X(G)}$ is

$$J_{X(G)} = \overline{\text{span}}\{fg^* : f, g \in C_c(G_1)\};$$

- (4) the Toeplitz representation (I_1, I_0) of Lemma 5 is Cuntz-Pimsner covariant.

Proof. (1) Choose $a \in C_r^*(G_0)$; by [6, Proposition II.4.2], $a \in C_0(G)$. Suppose that $a \notin \ker \phi$. Then there exists $\xi \in X(G)$ such that $a\xi = \phi(a)\xi$ is nonzero, so there exist composable γ, γ' with $a(\gamma) \neq 0$ and $\xi(\gamma') \neq 0$. We have $r(\gamma') \in r(G_1)$ by definition of $X(G)$, and so $a \notin C_0(\{g \in G_0 : s(g) \notin r(G_1)\})$.

On the other hand, suppose that $a \notin C_0(\{g \in G_0 : s(g) \notin r(G_1)\})$. Choose $g \in G_0$ such that $s(g) \in r(G_1)$ and $a(g) \neq 0$. Choose $\gamma \in G_1$ with $r(\gamma) = s(g)$, fix a precompact open bisection $U \subseteq G_1$ containing γ and use Urysohn's lemma to choose $\xi \in C_c(U) \subseteq C_c(G_1)$ such that $\xi(\gamma) = 1$. Then $(\phi(a)\xi)(g\gamma) = \sum_{\alpha\beta=g\gamma} a(\alpha)\xi(\beta)$. Since ξ is supported on the bisection U , and since $\alpha\beta = g\gamma$ implies $s(\beta) = s(\gamma)$, if $\alpha\beta = g\gamma$ and $a(\alpha)\xi(\beta) \neq 0$, we have $\beta = \gamma$ and then $\alpha = g$ by cancellation. So $(\phi(a)\xi)(g\gamma) = a(g) \neq 0$ and in particular $a \notin \ker \phi$.

(2) Suppose that $a \in \ker(\phi)^\perp$. Fix $g \in G_0$ with $s(g) \notin r(G_1)$. Since $G_1 = c^{-1}(1)$ is clopen and r is a local homeomorphism, $r(G_1)$ is also clopen and so there is an open $U \subseteq G^{(0)} \setminus r(G_1)$ with $s(g) \in U$. By Urysohn, there exists $f \in C_c(U)$ with $f(s(g)) = 1$. Part (1) gives $g \in \ker(\phi)$ and so $af = 0$, which gives $0 = (af)(g) = a(g)$. So $a \in C_0(\{g \in G_0 : s(g) \in r(G_1)\})$. On the other hand, if $a \in C_0(\{g \in G_0 : s(g) \in r(G_1)\})$ and $f \in \ker(\phi)$, then the preceding paragraph shows that $s(\{\gamma : a(\gamma) \neq 0\}) \cap r(\{\gamma : f(\gamma) \neq 0\}) = \emptyset$, and so $af = 0$ showing that $a \in \ker(\phi)^\perp$.

(3) We have seen that $r(G_1)$ is clopen, and since G_0 is also clopen, $s^{-1}(r(G_1)) \cap G_0$ is clopen. So for $a \in C_c(G_0)$ the pointwise products $a_1 = 1_{s^{-1}(r(G_1))}a$ and $a_0 = 1_{G_0 \setminus s^{-1}(r(G_1))}a$ belong to $C_c(G_0)$ and satisfy $a = a_0 + a_1$. Since $a_0 \in \ker(\phi)$, we have $\phi(a) = \phi(a_1)$. So $\phi(C_r^*(G_0)) = \overline{\phi(C_c(s^{-1}(r(G_1)) \cap G_0))}$. As observed above, if $\gamma \in s^{-1}(r(G_1)) \cap G_0$ then for any $\alpha \in G_1$ with $r(\alpha) = s(\gamma)$, we can write $\gamma = (\gamma\alpha)\alpha^{-1} \in G_1G_1^{-1}$. Since multiplication in G is open, it follows that

$$\{UV^{-1} : U, V \subseteq G_1 \text{ are precompact open bisections}\}$$

is a base for the topology on $s^{-1}(r(G_1)) \cap G_0$, and so $C_c(s^{-1}(r(G_1)) \cap G_0) = \overline{\text{span}}\{fg^* : f, g \in C_c(G_1)\}$. For $f, g \in C_c(G_1)$ and $\xi \in C_c(G_1) \subseteq X(G)$, we have $\phi(fg^*)(\xi) = fg^*\xi = \theta_{f,g}(\xi)$. So $\phi(fg^*) = \theta_{f,g} \in \mathcal{K}(X(G))$. Hence $\phi(C_c(s^{-1}(r(G_1)) \cap G_0)) \subseteq \mathcal{K}(X(G))$, and the result follows.

(4) Part (3) shows that $\text{span}\{fg^* : f, g \in C_c(G_1)\}$ is dense in the Katsura ideal $J_{X(G)}$. For $f, g \in C_c(G_1)$ we have $I_0(fg^*) = fg^*$ because I_0 extends the inclusion $C_c(G_0) \subseteq C_c(G)$ by definition, and we have just seen that $\phi(fg^*) = \theta_{f,g}$, giving $I_1^{(1)}(\phi(fg^*)) = I_1^{(1)}(\theta_{f,g}) = I_1(f)I_1(g)^* = fg^*$ as well. $\square$

Notation 9. Recall that if (ψ, π) is a Toeplitz representation of a C^* -correspondence X in a C^* -algebra B , then for $n \geq 2$ we write ψ_n for the continuous linear map from $X^{\otimes n}$ to B such that $\psi(x_1 \otimes \cdots \otimes x_n) = \psi(x_1) \cdots \psi(x_n)$ for all $x_i \in X$.

Proposition 10. *The integrated form $I_1 \times I_0$ of the representation in part (4) of Lemma 8 is an isomorphism of $\mathcal{O}_{X(G)}$ onto $C_r^*(G)$.*

Proof. The grading $c : C_r^*(G) \rightarrow \mathbb{Z}$ induces an action β of $\mathbb{T}$ on $C^*(G)$ such that

$$\beta_z(a) = z^{c(a)}a \quad \text{for all } z \in \mathbb{T} \text{ and } a \in \bigcup_n C_c(G_n).$$

Lemma 6 implies that for $n \geq 1$, the image $I_n(C_c(G_1)^{\otimes n})$ is dense in $C_c(G_n)$. This implies that $I_1 \times I_0$ is equivariant for the gauge action α on $\mathcal{O}_{X(G)}$ and β . The gauge-invariant uniqueness theorem [4, Theorem 6.4] implies that $I_1 \times I_0$ is injective. Moreover, since each $I_n(C_c(G_1)^{\otimes n})^*$ is dense in $C_c(G_n)^*$, which is $C_c(G_{-n})$ by Lemma 6, $I_1 \times I_0$ has dense range, and so is surjective since it is a homomorphism between C^* -algebras. $\square$

Corollary 11. *Suppose that $C^*(G_0) = C_r^*(G_0)$. Then $C^*(G) = C_r^*(G)$ and so Proposition 10 describes an isomorphism $\mathcal{O}_{X(G)} \cong C^*(G)$.*

Proof. We saw in Lemma 2 that the completion A_0 of $C_c(G_0)$ in $C_r^*(G)$ coincides with $C_r^*(G_0)$, and therefore, by hypothesis, with $C^*(G_0)$. So $\|I_0(a)\| = \|a\|_{C^*(G_0)}$ for all $a \in C_c(G_0)$. For $\xi \in C_c(G_1)$, we have

$$\|\xi^*\xi\|_{C^*(G)} \geq \|\xi^*\xi\|_{C_r^*(G)}.$$

The function $\xi^*\xi$ belongs to $C_c(G_0)$ and so $\|\xi^*\xi\|_{C_r^*(G)} = \|\xi^*\xi\|_{C_r^*(G_0)}$ by Lemma 2, and this last is equal to $\|\xi^*\xi\|_{C^*(G_0)}$ by hypothesis. Since the I -norm on $C_c(G)$ agrees with the I -norm on $C_c(G_0)$, the canonical inclusion $C_c(G_0) \hookrightarrow C_c(G)$ is I -norm bounded, and so extends to a C^* -homomorphism $C^*(G_0) \rightarrow C^*(G)$, which forces $\|\xi^*\xi\|_{C^*(G_0)} \geq \|\xi^*\xi\|_{C^*(G)}$. So we have equality throughout, giving

$$\|\xi^*\xi\|_{C^*(G)} = \|\xi^*\xi\|_{C_r^*(G_0)} = \|\langle I_1(\xi), I_1(\xi) \rangle_{C_r^*(G_0)}\|.$$

In particular, $\|\xi\|_{C^*(G)} = \|I_1(\xi)\|$. We deduce that (I_0, I_1) extend to a Cuntz-Pimsner covariant Toeplitz representation $(\tilde{I}_0, \tilde{I}_1)$ of $X(G)$ in $C^*(G)$. Let $q : C^*(G) \rightarrow C_r^*(G)$ denote the quotient map. Then $q \circ (\tilde{I}_1 \times \tilde{I}_0) = I_1 \times I_0$, which is injective by Proposition 10, and we deduce that q is also injective. $\square$

Remark.

If G_0 is an amenable groupoid (we don't have to specify which flavour because topological amenability and measurewise amenability coincide for second-countable locally compact Hausdorff étale groupoids [1, Theorem 3.3.7]), then [8, Proposition 9.3] shows that G is amenable as well, and then the preceding corollary follows. So the corollary has independent content only if G_0 is not amenable but nevertheless has identical full and reduced norms.

Let

$$L_0 = \{a \in M_2(C_c(G)) : a_{11} \in J_{X(G)}, a_{22} \in C_c(G_0), a_{12} \in C_c(G_1) \text{ and } a_{21} \in C_c(G_{-1})\}.$$

Then the completion of L_0 in the norm induced by the reduced norm on $C_r^*(G)$ is

$$L = \begin{pmatrix} J_{X(G)} & X(G) \\ X(G)^* & C_r^*(G_0) \end{pmatrix}.$$

Part (3) of Lemma 8 shows that L is a C^* -subalgebra of $M_2(C_r^*(G))$, and that the corners $\begin{pmatrix} J_{X(G)} & 0 \\ 0 & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 0 \\ 0 & C_r^*(G_0) \end{pmatrix}$ are full. So the inclusion maps $i^{11} : J_{X(G)} \rightarrow L$ into the top-left corner and $i^{22} : C_r^*(G_0) \rightarrow L$ into the bottom-right corner determine isomorphisms $i_*^{11} : K_*(J_{X(G)}) \rightarrow K_*(L)$ and $i_*^{22} : K_*(C_r^*(G_0)) \rightarrow K_*(L)$. The composite $[X] := (i_*^{22})^{-1} \circ i_*^{11} : K_*(J_{X(G)}) \rightarrow K_*(C_r^*(G_0))$ is the map appearing in [4, Theorem 8.6] (see [4, Remark B4]). The inclusion map $\iota : J_{X(G)} \hookrightarrow C_r^*(G_0)$ also induces a homomorphism $\iota_* : K_*(J_{X(G)}) \rightarrow K_*(C_r^*(G_0))$, and Theorem 8.6 of [4] gives the following.

Corollary 12. *There is an exact sequence in K -theory as follows:*

$$\begin{array}{ccccc} K_0(J_{X(G)}) & \xrightarrow{\iota_* - [X]} & K_0(C_r^*(G_0)) & \xrightarrow{(I_0)_*} & K_0(C_r^*(G)) \\ \uparrow & & & & \downarrow \\ K_1(C_r^*(G)) & \xleftarrow{(I_0)_*} & K_1(C_r^*(G_0)) & \xleftarrow{\iota_* - [X]} & K_1(J_{X(G)}) \end{array}$$

REFERENCES

- [1] C. Anantharaman-Delaroche and J. Renault, *Amenable groupoids*, With a foreword by Georges Skandalis and Appendix B by E. Germain, L'Enseignement Mathématique, Geneva, 2000, 196.
- [2] B. Blackadar, *Operator algebras: Theory of C^* -algebras and von Neumann algebras*, Encyclopaedia of Mathematical Sciences, **122**, Springer-Verlag, Berlin, 2006.
- [3] N. J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181.
- [4] T. Katsura, *On C^* -algebras associated to C^* -correspondences*, J. Funct. Anal. **217** (2004), 366–401.
- [5] M.V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by $\mathbf{Z}$* , Fields Inst. Commun., **12**, Free probability theory (Waterloo, ON, 1995), 189–212, Amer. Math. Soc., Providence, RI, 1997.
- [6] J. Renault, *A groupoid approach to C^* -algebras*, Lecture Notes in Mathematics **793** (1980).
- [7] A. Sims, B. Whitehead, and M. F. Whittaker, *Twisted C^* -algebras associated to finitely aligned higher-rank graphs*, preprint 2013 (arXiv:1310.7656 [math.OA]).
- [8] J. Spielberg, *Groupoids and C^* -algebras for categories of paths*, Trans. Amer. Math. Soc., to appear (arXiv:1111.6924v2 [math.OA]).

E-mail address: renniea@uow.edu.au, droberts@uow.edu.au, asims@uow.edu.au

SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOLLONGONG, WOLLONGONG, NSW, 2522, AUSTRALIA